



Institute for Advanced Laser Dentistry

Post-Op Packet

Post LANAP® Treatment Diet Instructions

- **The first three days** following Laser Therapy, follow **only** a liquid- like diet to allow healing. Anything that could be put into a blender to drink is ideal. The purpose of this is to protect the clot that is acting as a “band-aid” between the gums and the teeth. Do **not** drink through a straw, as this creates a vacuum in your mouth that can disturb the “band-aid”. Take daily vitamins.
- **Next four days after treatment,** foods with a “mushy” consistency such as those listed below are recommended.*see below
- **Starting seven to ten days after treatment,** soft foods may be allowable. The time to start on soft foods is dependent upon the loss of the white material that appeared around your teeth following LANAP® treatment. You were introduced to this material by your doctor or doctor’s assistant during your follow-up visit immediately following your initial treatment session. It was explained to you that you must leave this material alone until it naturally heals and disappears. Once the white material has disappeared, then soft foods can be introduced. Soft foods have the consistency of pasta, fish, chicken or steamed vegetables. You may then, gradually add back your regular diet choices.
- **Please remember** that even after ten days, healing is not complete. The first month following treatment you should continue to make smart food choices. Softer foods are better.

***“Mushy” Diet Suggestions**

DAILY VITAMINS!

Anything put through a food blender
Cream of wheat, Oatmeal, Malt O Meal
Mashed Avocado, Applesauce
Mashed Potatoes or Baked Potatoes – OK with butter/sour cream
Mashed Banana or any mashed/blended fruit except berries with seeds
Broth or Creamed Soup
Mashed steamed vegetables
Mashed Yams, Baked Sweet Potato or Butternut squash
Cottage cheese Cream or Soft cheese
Creamy peanut butter without solid pieces
Eggs any style, with or without melted cheese
Omelets can have cheese and avocado
Jell-O, Pudding, Ice Cream, Yogurt
Milk shake/smoothies – DO NOT blend with berries containing seeds
Ensure, Slim Fast -nutritional drinks

DON'T

Chew gum, candy, cookies, chips, nuts, anything hard or crunchy, anything that has seeds or hard pieces, meat that shreds and can lodge under the gum and between teeth, raw vegetables/salad.



Institute for Advanced Laser Dentistry

Post LANAP OHI "Loose Instructions"

Healing occurs at an individual's own rate and is affected by outside influence such as mechanical or bacterial trauma.

For example when you cut yourself and a scab forms, it will fall off as the tissue matures and becomes stronger and more resistant. How long does this process take for you, your father, your brother, your best friend and your son?

LOOSE OHI Guidelines:

- 1. manual brush at 7-10 days**
- 2. floss at 3-4 weeks (supra only)**
- 3. electric TB at 4-6 weeks (no sub-ging attachments)**
- 4. oral irrigators 3-6+ months (supra only)**
- 5. keep patients on the Chlorhexidine the entire time they are not brushing (about 2-4 weeks) and switch to a non-staining antimicrobial rinse indefinitely as preventative OHI.**

If there is fibrin or red thrombus still present or if the tissue does not look right, they can not brush yet. The tissue needs to be mature and strong enough for flossing and electric TB. Be the physician and diagnose before you make your OHI recommendations.

If you are experiencing severe pain, increasing swelling or uncontrollable bleeding, please call the office. After office hours, you can reach Dr. Lesco at home 410-560-2151 or cell phone 410-935-4282.